



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 990 437 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
05.04.2000 Bulletin 2000/14

(51) Int. Cl.⁷: **A61K 9/00, A61K 9/12,
A61K 31/56, A61K 31/135**

(21) Application number: **99204248.1**

(22) Date of filing: **04.12.1992**

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL
PT SE**

(30) Priority: **12.12.1991 GB 9126378
12.12.1991 GB 9126405
06.02.1992 GB 9202522**

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
**96202592.0 / 0 756 868
92924667.6 / 0 616 523**

(71) Applicant: **GLAXO GROUP LIMITED
Greenford, Middlesex UB6 0NN (GB)**

(72) Inventors:
• **Marriott, Rachel Ann
San Diego, CA 92129 (US)**
• **Taylor, Anthony James
Ware, Hertfordshire SG12 0DP (GB)**
• **Wyatt, David Andrew
Ware, Hertfordshire SG12 0DP (GB)**

(74) Representative:
**Teuten, Andrew John et al
Glaxo Wellcome plc,
Glaxo Wellcome House,
Berkeley Avenue
Greenford, Middlesex UB6 0NN (GB)**

(54) **Aerosol compositions**

(57) This invention relates to aerosol formulations of use for the administration of medicaments by inhalation, in particular a pharmaceutical aerosol formulation which comprises particulate medicament selected from the group comprising salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates thereof and a fluorocarbon or hydrogen-containing chlorofluorocarbon propellant, which formulation is substantially free of surfactant.

A method of treating respiratory disorders which comprises administration by inhalation of an effective amount of a pharmaceutical aerosol formulation as defined is also described.

EP 0 990 437 A1

Description

[0001] This invention relates to aerosol formulations of use for the administration of medicaments by inhalation.

[0002] The use of aerosols to administer medicaments has been known for several decades. Such aerosols generally comprise the medicament, one or more chlorofluorocarbon propellants and either a surfactant or a solvent, such as ethanol. The most commonly used aerosol propellants for medicaments have been propellant 11 (CCl_3F) and/or propellant 114 ($\text{CF}_2\text{ClCF}_2\text{Cl}$) with propellant 12 (CCl_2F_2). However these propellants are now believed to provoke the degradation of stratospheric ozone and there is thus a need to provide aerosol formulations for medicaments which employ so called "ozone-friendly" propellants.

[0003] A class of propellants which are believed to have minimal ozone-depleting effects in comparison to conventional chlorofluorocarbons comprise fluorocarbons and hydrogen-containing chlorofluorocarbons, and a number of medicinal aerosol formulations using such propellant systems are disclosed in, for example, EP 0372777, WO91/04011, WO91/11173, WO91/11495 and WO91/14422. These applications are all concerned with the preparation of pressurised aerosols for the administration of medicaments and seek to overcome the problems associated with the use of the new class of propellants, in particular the problems of stability associated with the pharmaceutical formulations prepared. The applications all propose the addition of one or more of adjuvants such as alcohols, alkanes, dimethyl ether, surfactants (including fluorinated and non-fluorinated surfactants, carboxylic acids, polyethoxylates etc) and even conventional chlorofluorocarbon propellants in small amounts intended to minimise potential ozone damage.

[0004] Thus, for example EP 0372777 requires the use of 1,1,1,2-tetrafluoroethane in combination with both a cosolvent having greater polarity than 1,1,1,2-tetrafluoroethane (e.g. an alcohol or a lower alkane) and a surfactant in order to achieve a stable formulation of a medicament powder. In particular it is noted in the specification at page 3, line 7 that "it has been found that the use of propellant 134a (1,1,1,2-tetrafluoroethane) and drug as a binary mixture or in combination with a conventional surfactant such as sorbitan trioleate does not provide formulations having suitable properties for use with pressurised inhalers". Surfactants are generally recognised by those skilled in the art to be essential components of aerosol formulations, required not only to reduce aggregation of the medicament but also to lubricate the valve employed, thereby ensuring consistent reproducibility of valve actuation and accuracy of dose dispensed. Whilst WO91/11173, WO91/11495 and WO91/14422 are concerned with formulations comprising an admixture of drug and surfactant, WO91/04011 discloses medicinal aerosol formulations in which the particulate medicaments are pre-coated with surfactant prior to dispersal in 1,1,1,2-tetrafluoroethane.

[0005] We have now surprisingly found that, in contradistinction to these teachings, it is in fact possible to obtain satisfactory dispersions of certain medicaments in fluorocarbon or hydrogen-containing chlorofluorocarbon propellants such as 1,1,1,2-tetrafluoroethane without recourse to the use of any surfactant or cosolvent in the composition, or the necessity to pre-treat the medicament prior to dispersal in the propellant. More particularly, satisfactory dispersions may be formed where the medicament is selected from salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates thereof.

[0006] There is thus provided in one aspect of the invention a pharmaceutical aerosol formulation which comprises particulate medicament selected from the group consisting of salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates (for example hydrates) thereof and a fluorocarbon or hydrogen-containing chlorofluorocarbon propellant, which formulation is substantially free of surfactant. By "substantially free of surfactant" is meant formulations which contain no significant amounts of surfactant, for example less than 0.0001% by weight of the medicament.

[0007] In an alternative embodiment the present invention provides a pharmaceutical aerosol formulation as hereinbefore defined with the proviso that when said formulation consists essentially of salbutamol and 1,1,1,2-tetrafluoroethane in a weight ratio of 0.05:18, said salbutamol is present in the form of a physiologically acceptable salt.

[0008] The particle size of the particulate (e.g. micronised) medicament should be such as to permit inhalation of substantially all of the medicament into the lungs upon administration of the aerosol formulation and will thus be less than 100 microns, desirably less than 20 microns, and preferably in the range 1-10 microns, e.g. 1-5 microns.

[0009] Suitable pharmaceutically acceptable salts of the medicaments of use in the formulations of the present invention include acid addition salts such as for example sulphates, hydrochlorides and xinafoates (1-hydroxy-2-naphthoate), amine salts or alkali metal salts (e.g. sodium). Salmeterol will preferably be in the form of its xinafoate salt and salbutamol will preferably be in the form of its sulphate salt.

[0010] The final aerosol formulation desirably contains 0.005-10% w/w, preferably 0.005 - 5% w/w, especially 0.01-1.0% w/w, of medicament relative to the total weight of the formulation.

[0011] The propellants for use in the invention may be any fluorocarbon or hydrogen-containing chlorofluorocarbon or mixtures thereof having a sufficient vapour pressure to render them effective as propellants. Preferably the propellant will be a non-solvent for the medicament. Suitable propellants include, for example, C_{1-4} hydrogen-containing chlorofluorocarbons such as CH_2ClF , $\text{CClF}_2\text{CHClF}$, CF_3CHClF , $\text{CHF}_2\text{CClF}_2$, CHClFCHF_2 , $\text{CF}_3\text{CH}_2\text{Cl}$ and CClF_2CH_3 ; C_{1-4} hydrogen-containing fluorocarbons such as CHF_2CHF_2 , $\text{CF}_3\text{CH}_2\text{F}$, CHF_2CH_3 and CF_3CHF_2 ; and perfluorocarbons

such as CF_3CF_3 and $\text{CF}_3\text{CF}_2\text{CF}_3$.

[0012] Where mixtures of the fluorocarbons or hydrogen-containing chlorofluorocarbons are employed they may be mixtures of the above identified compounds or mixtures, preferably binary mixtures, with other fluorocarbons or hydrogen-containing chlorofluorocarbons for example CHClF_2 , CH_2F_2 and CF_3CH_3 . Preferably a single fluorocarbon or hydrogen-containing chlorofluorocarbon is employed as the propellant. Particularly preferred as propellants are C_1 - C_4 hydrogen-containing fluorocarbons such as 1,1,1,2-tetrafluoroethane ($\text{CF}_3\text{CH}_2\text{F}$) and 1,1,1,2,3,3,3-heptafluoro-n-propane ($\text{CF}_3\text{CHFCF}_3$).

[0013] It is desirable that the formulations of the invention contain no components which may provoke the degradation of stratospheric ozone. In particular it is desirable that the formulations are substantially free of chlorofluorocarbons such as CCl_3F , CCl_2F_2 and CF_3CCl_3 .

[0014] The propellant may additionally contain a volatile adjuvant such as a saturated hydrocarbon for example propane, n-butane, isobutane, pentane and isopentane or a dialkyl ether for example dimethyl ether. In general, up to 50% w/w of the propellant may comprise a volatile hydrocarbon, for example 1 to 30% w/w. However, formulations which are substantially free of volatile adjuvants are preferred.

[0015] It is further desirable that the formulations of the invention are substantially free of liquid components of higher polarity than the propellant employed. Polarity may be determined for example, by the method described in European Patent Application Publication No. 0327777. In particular formulations which are substantially free of alcohols such as ethanol are preferable. As used herein "substantially free" means less than 1% w/w based upon the fluorocarbon or hydrogen-containing chlorofluorocarbon, in particular less than 0.5% for example 0.1% or less.

[0016] A particularly preferred embodiment of the invention provides a pharmaceutical aerosol formulation consisting essentially of one or more particulate medicament selected from the group consisting of salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates thereof, and one or more fluorocarbon or hydrogen-containing chlorofluorocarbon propellant.

[0017] It will be appreciated by those skilled in the art that the aerosol formulations according to the invention may, if desired, contain a combination of two or more active ingredients. Aerosol compositions containing two active ingredients (in a conventional propellant system) are known, for example, for the treatment of respiratory disorders such as asthma. Accordingly the present invention further provides aerosol formulations in accordance with the invention which contain two or more particulate medicaments. Medicaments may be selected from suitable combinations of the medicaments mentioned hereinbefore or may be selected from any other suitable drug useful in inhalation therapy and which may be presented in a form which is substantially completely insoluble in the selected propellant. Appropriate medicaments may thus be selected from, for example, analgesics, e.g. codeine, dihydromorphine, ergotamine, fentanyl or morphine; anginal preparations, e.g. diltiazem; antiallergics, e.g. cromoglycate, ketotifen or nedocromil; anti-infectives e.g. cephalosporins, penicillins, streptomycin, sulphonamides, tetracyclines and pentamidine; antihistamines, e.g. methapyrilene; anti-inflammatories, e.g. flunisolide, budesonide, tiopredane or triamcinolone acetonide; antitussives, e.g. noscapine; bronchodilators, e.g. ephedrine, adrenaline, fenoterol, formoterol, isoprenaline, metaproterenol, phenylephrine, phenylpropanolamine, pirbuterol, reproterol, rimeterol, terbutaline, isoetharine, tulobuterol, orciprenaline, or (-)-4-amino-3,5-dichloro- α -[[[6-(2-(2-pyridinyl)ethoxy)hexyl]amino]methyl]benzenemethanol; diuretics, e.g. amiloride; anticholinergics e.g. ipratropium, atropine or oxitropium; hormones, e.g. cortisone, hydrocortisone or prednisolone; xanthines e.g. aminophylline, choline theophyllinate, lysine theophyllinate or theophylline; and therapeutic proteins and peptides, e.g. insulin or glucagon. It will be clear to a person skilled in the art that, where appropriate, the medicaments may be used in the form of salts (e.g. as alkali metal or amine salts or as acid addition salts) or as esters (e.g. lower alkyl esters) or as solvates (e.g. hydrates) to optimise the activity and/or stability of the medicament and/or to minimise the solubility of the medicament in the propellant.

[0018] Particularly preferred aerosol formulations contain salbutamol (e.g. as the free base or the sulphate salt) or salmeterol (e.g. as the xinafoate salt) in combination with an anti-inflammatory steroid such as a beclomethasone ester (e.g. the dipropionate) or a fluticasone ester (e.g. the propionate) or an antiallergic such as cromoglycate (e.g. the sodium salt). Combinations of salmeterol and fluticasone propionate or beclomethasone dipropionate, or salbutamol and fluticasone propionate or beclomethasone dipropionate are preferred, especially salmeterol xinafoate and fluticasone propionate or salbutamol and beclomethasone dipropionate.

[0019] The formulations of the invention may be prepared by dispersal of the medicament in the selected propellant in an appropriate container, e.g. with the aid of sonication. The process is desirably carried out under anhydrous conditions to obviate any adverse effects of moisture on suspension stability.

[0020] The formulations according to the invention form weakly flocculated suspensions on standing but, surprisingly, these suspensions have been found to be easily redispersed by mild agitation to provide suspensions with excellent delivery characteristics suitable for use in pressurised inhalers, even after prolonged storage. Minimising and preferably avoiding the use of formulation excipients e.g. surfactants, cosolvents etc in the aerosol formulations according to the invention is also advantageous since the formulations may be substantially taste and odour free, less irritant and less toxic than conventional formulations.

[0021] The chemical and physical stability and the pharmaceutical acceptability of the aerosol formulations according to the invention may be determined by techniques well known to those skilled in the art. Thus, for example, the chemical stability of the components may be determined by HPLC assay, for example, after prolonged storage of the product. Physical stability data may be gained from other conventional analytical techniques such as, for example, by leak testing, by valve delivery assay (average shot weights per actuation), by dose reproducibility assay (active ingredient per actuation) and spray distribution analysis.

[0022] The particle size distribution of the aerosol formulations according to the invention is particularly impressive and may be measured by conventional techniques, for example by cascade impaction or by the "Twin Impinger" analytical process. As used herein reference to the "Twin Impinger" assay means "Determination of the deposition of the emitted dose in pressurised inhalations using apparatus A" as defined in British Pharmacopoeia 1988, pages A204-207, Appendix XVII C. Such techniques enable the "respirable fraction" of the aerosol formulations to be calculated. As used herein reference to "respirable fraction" means the amount of active ingredient collected in the lower impingement chamber per actuation expressed as a percentage of the total amount of active ingredient delivered per actuation using the twin impinger method described above. The formulations according to the invention have been found to have a respirable fraction of 20% or more by weight of the medicament, preferably 25 to 70%, for example 30 to 60%.

[0023] Optionally, the medicament may be surface-modified prior to its dispersion in the propellant by treatment with a substantially non-polar liquid medium which is a non-solvent for the medicament. There is thus provided in a further aspect of the invention an aerosol formulation comprising particulate, surface-modified medicament, as defined herein, and a fluorocarbon or hydrogen-containing chlorofluorocarbon propellant, which formulation is substantially free of surfactant. By "surface-modified medicament" is meant particles of medicament selected from the group consisting of salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates thereof which have been surface-modified by admixture with a substantially non-polar non-solvent liquid, followed by removal of the liquid. The substantially non-polar non-solvent liquid medium is conveniently an aliphatic hydrocarbon, e.g. a lower alkane, which is sufficiently volatile to permit its ready evaporation, e.g. at ambient temperature and pressure, after slurring with the medicament. The use of isopentane as liquid medium is particularly advantageous in this respect.

[0024] The medicament is desirably slurried with the liquid medium under anhydrous conditions to obviate any adverse effects of moisture on suspension stability. The slurry may advantageously be sonicated to maximise the surface-modifying effect of the treatment. The liquid may be removed by any convenient means for example by evaporation or by filtration followed by evaporation, provided that following treatment the medicament is substantially free of the liquid. The formulations of the invention will be substantially free of the non-solvent non-polar liquid. Surface-modified medicament prepared by the above-described process comprises a further aspect of the present invention.

[0025] The formulations according to the invention may be filled into canisters suitable for delivering pharmaceutical aerosol formulations. Canisters generally comprise a container capable of withstanding the vapour pressure of the propellant used such as a plastic or plastic-coated glass bottle or preferably a metal can, for example an aluminium can which may optionally be anodised, lacquer-coated and/or plastic-coated, which container is closed with a metering valve. The metering valves are designed to deliver a metered amount of the formulation per actuation and incorporate a gasket to prevent leakage of propellant through the valve. The gasket may comprise any suitable elastomeric material such as for example low density polyethylene, chlorobutyl, black and white butadiene-acrylonitrile rubbers, butyl rubber and neoprene. Suitable valves are commercially available from manufacturers well known in the aerosol industry, for example, from Valois, France (e.g. DF10, DF30, DF60), Bepak plc, UK (e.g. BK300, BK356) and 3M-Neotech Ltd, UK (e.g. Spraymiser™).

[0026] Conventional bulk manufacturing methods and machinery well known to those skilled in the art of pharmaceutical aerosol manufacture may be employed for the preparation of large scale batches for the commercial production of filled canisters. Thus, for example, in one bulk manufacturing method a metering valve is crimped onto an aluminium can to form an empty canister. The particulate medicament is added to a charge vessel and liquified propellant is pressure filled through the charge vessel into a manufacturing vessel. The drug suspension is mixed before recirculation to a filling machine and an aliquot of the drug suspension is then filled through the metering valve into the canister. Typically, in batches prepared for pharmaceutical use, each filled canister is check-weighed, coded with a batch number and packed into a tray for storage before release testing.

[0027] Each filled canister is conveniently fitted into a suitable channelling device prior to use to form a metered dose inhaler for administration of the medicament into the lungs or nasal cavity of a patient. Suitable channelling devices comprise for example a valve actuator and a cylindrical or cone-like passage through which medicament may be delivered from the filled canister via the metering valve to the nose or mouth of a patient e.g. a mouthpiece actuator. Metered dose inhalers are designed to deliver a fixed unit dosage of medicament per actuation or "puff", for example in the range of 10 to 5000 microgram medicament per puff.

[0028] Administration of medicament may be indicated for the treatment of mild, moderate or severe acute or chronic symptoms or for prophylactic treatment. It will be appreciated that the precise dose administered will depend on the age

and condition of the patient, the particular particulate medicament used and the frequency of administration and will ultimately be at the discretion of the attendant physician. When combinations of medicaments are employed the dose of each component of the combination will in general be that employed for each component when used alone. Typically, administration may be one or more times, for example from 1 to 8 times per day, giving for example 1,2,3 or 4 puffs each time.

[0029] Suitable daily doses, may be, for example in the range 50 to 200 microgram of salmeterol, 100 to 1000 microgram of salbutamol, 50 to 2000 microgram of fluticasone propionate or 100 to 2000 microgram of beclomethasone dipropionate, depending on the severity of the disease.

[0030] Thus, for example, each valve actuation may deliver 25 microgram salmeterol, 100 microgram salbutamol, 25, 50, 125 or 250 microgram fluticasone propionate or 50, 100, 200 or 250 microgram beclomethasone dipropionate. Typically each filled canister for use in a metered dose inhaler contains 100, 160 or 240 metered doses or puffs of medicament.

[0031] The filled canisters and metered dose inhalers described herein comprise further aspects of the present invention.

[0032] A still further aspect of the present invention comprises a method of treating respiratory disorders such as, for example, asthma, which comprises administration by inhalation of an effective amount of a formulation as herein described.

[0033] The following non-limitative Examples serve to illustrate the invention.

Example 1

[0034] Micronised salmeterol xinafoate (24mg) was weighed into a clean, dry, plastic-coated glass bottle and 1,1,1,2-tetrafluoroethane (18.2g) was added from a vacuum flask. The bottle was quickly sealed with a blank aluminium ferrule. The resulting aerosol contained 0.132% w/w salmeterol xinafoate.

Example 2

[0035] Micronised salmeterol xinafoate (38.28g) and 1,1,1,2-tetrafluoroethane (36.36kg) were added to a pressure vessel and mixed with a high shear mixer for 20 minutes. Aliquots (18.2g) of the suspension were filled into aluminium cans closed with a metering valve, filling under pressure through the valve using conventional filling equipment. The resulting inhalers contained 9.57mg salmeterol xinafoate and delivered 25 microgram salmeterol (39.9 microgram salt) per actuation.

Example 3

[0036] Micronised fluticasone propionate (24mg) was weighed into a clean, dry, plastic-coated glass bottle and 1,1,1,2-tetrafluoroethane (18.2g) was added from a vacuum flask. The bottle was quickly sealed with a blank aluminium ferrule. The resulting aerosol contained 0.132% w/w fluticasone propionate.

Examples 4 and 5

[0037] Micronised fluticasone propionate (66mg or 6.6mg) was weighed directly into each of 100 open aluminium cans and a metering valve was then crimped into place on each can. 1,1,1,2-Tetrafluoroethane (18.2g) was then added to each canister under pressure, through the valve, and each filled canister shaken to disperse the drug. The resulting inhalers contained 66 or 6.6mg fluticasone propionate and delivered 250 or 25 microgram fluticasone propionate per actuation (Examples 4 and 5 respectively).

Example 6

[0038] Micronised salbutamol (24mg) was weighed into a clean, dry, plastic-coated glass bottle and 1,1,1,2-tetrafluoroethane (18.2g) was added from a vacuum flask. The bottle was quickly sealed with a blank aluminium ferrule. The resulting aerosol contained 0.132% w/w salbutamol.

Examples 7 and 8

[0039] Micronised salbutamol (24mg or 48mg) was weighed directly into each of 3 open aluminium cans. 1,1,1,2-Tetrafluoroethane (18.2g) was added to each can from a vacuum flask and a metering valve was then crimped into place. Each filled canister was then shaken in an ultrasonic bath for 8 minutes. The resulting inhalers contained 24mg

or 48mg salbutamol and delivered 100 or 200 microgram salbutamol per actuation (Examples 7 and 8 respectively).

Example 9

5 [0040] Micronised salbutamol sulphate (31.7mg) was weighed into a clean, dry, plastic-coated glass bottle and 1,1,1,2-tetrafluoroethane (18.2g) was added from a vacuum flask. The bottle was quickly sealed with a blank aluminium ferrule. The resulting aerosol contained 0.174% w/w salbutamol sulphate.

Example 10

10 [0041] Micronised salbutamol sulphate (31.7mg) was weighed directly into each of 4 open aluminium cans. 1,1,1,2-Tetrafluoroethane (18.2g) was added to each can from a vacuum flask and a metering valve was then crimped into place. Each filled canister was then shaken in an ultrasonic bath for 5 minutes. The resulting inhalers contained 31.7mg salbutamol sulphate and delivered 100 microgram salbutamol per actuation.

Example 11

15 [0042] Isopentane (25ml) was added to micronised salmeterol xinafoate (0.5g) to form a slurry, which was sonicated for 3 minutes. The resulting suspension was dried by evaporating the isopentane at ambient temperature to yield surface-modified salmeterol xinafoate. Samples of this product (11.6mg) were weighed into aluminium aerosol cans and 1,1,1,2-tetrafluoroethane (18.2g - 99.95% w/w of total fill weight) was added to each can, whereafter suitable metering valves were crimped onto the cans, which were then each sonicated for 5 minutes. The resulting aerosols contained salmeterol in an amount equivalent to 240 actuations at 25 microgram per actuation.

25 Example 12

[0043] Micronised beclomethasone dipropionate monohydrate (68 mg) was weighed into a clean, dry, plastic-coated glass bottle and 1,1,1,2-tetrafluoroethane (to 18.2g) was added from a vacuum flask. The bottle was quickly sealed with a metering valve. The resulting aerosol dispensed 250 microgram beclomethasone dipropionate (as the monohydrate) per 75.8mg actuation.

Example 13

30 [0044] Micronised salmeterol xinafoate (9.57mg) is weighed directly into an aluminium can and 1,1,1,2,3,3,3-heptafluoro-n-propane (to 21.4g) added from a vacuum flask. A metering valve is crimped into place and the filled canister sonicated for five minutes. The aerosol delivers 25 microgram salmeterol per actuation.

Example 14

40 [0045] Micronised fluticasone propionate (13.3mg) is weighed directly into an aluminium can and 1,1,1,2,3,3,3-heptafluoro-n-propane (to 21.4g) added from a vacuum flask. A metering valve is crimped into place and the filled canister sonicated for five minutes. The aerosol delivers 50 microgram fluticasone propionate per actuation.

Example 15

45 [0046] Micronised salbutamol sulphate (29mg) was weighed directly into an aluminium can and 1,1,1,2,3,3,3-heptafluoro-n-propane (to 21.4g) added from a vacuum flask. A metering valve was crimped into place and the filled canister sonicated for five minutes. The aerosol delivered 100 microgram salbutamol per actuation.

50 Example 16

[0047] Micronised beclomethasone dipropionate monohydrate (62mg) was weighed directly into an aluminium can and 1,1,1,2,3,3,3-heptafluoro-n-propane (to 21.4g) added from a vacuum flask. A metering valve was crimped into place and the filled canister sonicated for five minutes. The aerosol delivered 250 microgram beclomethasone dipropionate per actuation.

Example 17**[0048]**

5

10

	Per Inhaler % w/w	Per Actuation
Salmeterol xinafoate	0.048	36.25 microgram
Fluticasone propionate	0.066	50 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8mg

15 **[0049]** Micronised medicaments were weighed into an aluminium can, 1,1,1,2-tetrafluoroethane (18.2g) was added from a vacuum flask and a metering valve was crimped into place.

Example 18**[0050]**

20

25

	Per Inhaler % w/w	Per Actuation
Salmeterol xinafoate	0.048	36.25 microgram
Fluticasone propionate	0.165	125 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8mg

30 **[0051]** Micronised medicaments were weighed into an aluminium can, 1,1,1,2-tetrafluoroethane (18.2g) was added from a vacuum flask and a metering valve was crimped into place.

Example 1935 **[0052]**

40

	Per Inhaler % w/w	Per Actuation
Salmeterol xinafoate	0.048	36.25 microgram
Fluticasone propionate	0.132	100 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8mg

45

Example 20**[0053]**

50

55

	Per Inhaler % w/w	Per Actuation
Salmeterol xinafoate	0.048	36.25 microgram
Fluticasone propionate	0.330	250 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8mg

Example 21

[0054]

	Per Inhaler % w/w	Per Actuation
Salbutamol *	0.132	100 microgram
Fluticasone propionate	0.132	100 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8mg

* as free base or an equivalent weight of salt e.g. sulphate

Example 22

[0055]

	Per Inhaler % w/w	Per Actuation
Salbutamol *	0.264	200 microgram
Fluticasone propionate	0.330	250 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8mg

* as free base or an equivalent weight of salt e.g. sulphate

Example 23

[0056]

	Per Inhaler % w/w	Per Actuation
Salmeterol xinafoate	0.048	36.25 microgram
Beclomethasone dipropionate	0.066	50 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8mg

Example 24

[0057]

	Per Inhaler % w/w	Per Actuation
Salmeterol xinafoate	0.048	36.25 microgram
Fluticasone propionate	0.264	200 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8mg

Example 25**[0058]**

5

10

	Per Inhaler % w/w	Per Actuation
Salbutamol *	0.132	100 microgram
Beclomethasone dipropionate	0.066	50 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8mg

* as free base or an equivalent weight of salt e.g. sulphate

15

Example 26**[0059]**

20

25

	Per Inhaler % w/w	Per Actuation
Salbutamol *	0.264	200 microgram
Beclomethasone dipropionate	0.264	200 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8mg

* as free base or an equivalent weight of salt e.g. sulphate

30

[0060] In Examples 19 to 26 micronised medicaments are weighed into aluminium cans, 1,1,1,2-tetrafluoroethane (18.2g) is added from a vacuum flask, and metering valves are crimped into place.

Claims

35

40

45

50

55

1. A pharmaceutical aerosol formulation which comprises particulate medicament selected from the group consisting of salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates thereof and a fluorocarbon or hydrogen-containing chlorofluorocarbon propellant, which formulation is substantially free of surfactant.
2. A pharmaceutical aerosol formulation which comprises particulate medicament selected from the group consisting of salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates thereof and a fluorocarbon or hydrogen-containing chlorofluorocarbon propellant, which formulation is substantially free of surfactant and with the proviso that when said formulation consists essentially of salbutamol and 1,1,1,2-tetrafluoroethane in a weight ratio of 0.05:18, said salbutamol is present in the form of a physiologically acceptable salt.
3. A pharmaceutical aerosol formulation consisting essentially of one or more particulate medicament selected from the group consisting of salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates thereof, and one or more fluorocarbon or hydrogen-containing chlorofluorocarbon propellant.
4. A formulation as claimed in any one of claims 1 to 3 wherein the medicament is salmeterol xinafoate.
5. A formulation as claimed in any one of claims 1 to 3 wherein the medicament is salbutamol sulphate.
6. A formulation as claimed in any one of claims 1 to 3 wherein the medicament is fluticasone propionate.

7. A formulation as claimed in any one of claims 1 to 3 wherein the medicament is beclomethasone dipropionate or a physiologically acceptable solvate thereof.
8. A formulation as claimed in one of claims 1 to 7 wherein the propellant comprises 1,1,1,2-tetrafluoroethane or 1,1,1,2,3,3,3-heptafluoro-n-propane.
9. A formulation as claimed in one of claims 1 to 8 wherein the propellant comprises 1,1,1,2-tetrafluoroethane.
10. A formulation as claimed in any one of claims 1 to 9 wherein the medicament is present in an amount of 0.005 to 10% w/w based on the total weight of the formulation.
11. A formulation as claimed in any one of claims 1 to 10 which contains two or more particulate medicaments.
12. A formulation as claimed in Claim 11 which contains salbutamol or salmeterol or a physiologically acceptable salt thereof in combination with an antiinflammatory steroid or an antiallergic.
13. A formulation as claimed in Claim 12 which contains salmeterol or salbutamol or a physiologically acceptable salt thereof in combination with fluticasone propionate or beclomethasone dipropionate or a physiologically acceptable solvate thereof.
14. A formulation as claimed in Claim 13 which contains salmeterol xinafoate and fluticasone propionate.
15. A formulation as claimed in Claim 13 which contains salbutamol and beclomethasone dipropionate.
16. A formulation as claimed in any one of claims 1 to 15 which has a respirable fraction of 20% or more by weight of the medicament.
17. A formulation as claimed in any one of claims 1 to 16 wherein said particulate medicament is surface-modified.
18. Surface-modified medicament prepared by admixture of particles of a medicament selected from the group consisting of salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates thereof, with a substantially non-polar, non-solvent liquid, followed by removal of the liquid.
19. A canister suitable for delivering a pharmaceutical aerosol formulation which comprises a container capable of withstanding the vapour pressure of the propellant used, which container is closed with a metering valve and contains a pharmaceutical aerosol formulation which comprises particulate medicament selected from the group consisting of salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates thereof and a fluorocarbon or hydrogen-containing chlorofluorocarbon propellant, which formulation is substantially free of surfactant.
20. A canister as claimed in claim 19 wherein the container is a metal can.
21. A metered dose inhaler which comprises a canister as claimed in claim 19 or claim 20 fitted into a suitable channelling device.
22. A method of treating respiratory disorders which comprises administration by inhalation of an effective amount of a pharmaceutical aerosol formulation which comprises particulate medicament selected from the group consisting of salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates thereof and a fluorocarbon or hydrogen-containing chlorofluorocarbon propellant, which formulation is substantially free of surfactant.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 20 4248

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A,D	WO 91 11173 A (FISONS) 8 August 1991 (1991-08-08) * claims * * page 3, line 15 - line 16 * * page 7, line 19 - line 20 * * page 8, line 1 - line 7 *	1-22	A61K9/00 A61K9/12 A61K31/56 A61K31/135
A	WO 91 11496 A (BOEHRINGER INGELHEIM) 8 August 1991 (1991-08-08) * claims *	1-22	
A,P	WO 92 08447 A (GLAXO) 29 May 1992 (1992-05-29) * claims *	1-22	
A,P	WO 92 00061 A (FISONS) 9 January 1992 (1992-01-09) * claims * * examples * * page 7, line 1 - line 6 *	1-22	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			A61K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 9 February 2000	Examiner Scarpont, U
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			

EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 20 4248

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

09-02-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9111173 A	08-08-1991	AT 125153 T	15-08-1995
		CA 2074495 A	03-08-1991
		DE 69111426 D	24-08-1995
		DE 69111426 T	18-01-1996
		DK 513127 T	30-10-1995
		EP 0513127 A	19-11-1992
		ES 2075956 T	16-10-1995
		GR 3017611 T	31-01-1996
		IE 67185 B	06-03-1996
		IL 97065 A	25-01-1994
		JP 2858948 B	17-02-1999
		JP 5503523 T	10-06-1993
		MX 24390 A	01-04-1993
		NZ 236948 A	26-08-1992
		PT 96639 A,B	31-10-1991
		ZA 9100696 A	30-10-1991
WO 9111496 A	08-08-1991	DE 4003270 A	08-08-1991
		AT 185587 T	15-10-1999
		AU 656129 B	27-01-1995
		AU 7211691 A	21-08-1991
		CA 2075060 A	04-08-1991
		CS 9100265 A	15-09-1991
		DE 59109161 D	18-11-1999
		EP 0513099 A	19-11-1992
		FI 923491 A	03-08-1992
		HR 940737 A	30-06-1997
		JP 5504350 T	08-07-1993
		NO 302420 B	02-03-1998
		NZ 236974 A	27-07-1993
		PT 96635 A,B	31-10-1991
		SI 9110155 A,B	30-04-1998
		ZA 9100756 A	28-10-1992
WO 9208447 A	29-05-1992	AT 127012 T	15-09-1995
		AU 653369 B	29-09-1994
		AU 8862991 A	11-06-1992
		CA 2094727 A	10-05-1992
		DE 69112635 D	05-10-1995
		DE 69112635 T	08-02-1996
		DK 556239 T	06-11-1995
		EP 0556239 A	25-08-1993
		ES 2078550 T	16-12-1995
		GR 3018171 T	29-02-1996
		JP 6501693 T	24-02-1994
		US 5785952 A	28-07-1998

EPO FORM P4459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 20 4248

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

09-02-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9200061 A	09-01-1992	AT 147973 T	15-02-1997
		AU 8055691 A	23-01-1992
		CA 2085884 A	30-12-1991
		DE 69124374 D	06-03-1997
		DE 69124374 T	19-06-1997
		DK 536235 T	14-04-1997
		EP 0536235 A	14-04-1993
		ES 2096653 T	16-03-1997
		FI 925852 A	23-12-1992
		GR 3023065 T	30-07-1997
		IL 98666 A	12-04-1994
		JP 2854974 B	10-02-1999
		MX 9100008 A	03-02-1992
		NO 924954 A	21-12-1992
		NZ 238746 A	28-10-1992
		PT 98133 A	30-04-1992
		SK 392592 A	06-07-1994
		US 5846521 A	08-12-1998
		ZA 9104897 A	25-03-1992

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82



(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
20.11.2002 Bulletin 2002/47

(51) Int Cl.7: **A61K 9/00, A61K 9/12,**
A61K 31/56, A61K 31/135

(21) Application number: **99204248.1**

(22) Date of filing: **04.12.1992**

(54) **Aerosol compositions**

Aerosolpräparate

Compositions pharmaceutiques sous forme d'aérosol

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL
PT SE

(30) Priority: **12.12.1991 GB 9126378**
12.12.1991 GB 9126405
06.02.1992 GB 9202522

(43) Date of publication of application:
05.04.2000 Bulletin 2000/14

(60) Divisional application:
00202961.9 / 1 066 828

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
96202592.0 / 0 756 868
92924667.6 / 0 616 523

(73) Proprietor: **GLAXO GROUP LIMITED**
Greenford, Middlesex UB6 ONN (GB)

(72) Inventors:
• **Marriott, Rachel Ann**
San Diego, CA 92129 (US)
• **Taylor, Anthony James**
Ware, Hertfordshire SG12 0DP (GB)
• **Wyatt, David Andrew**
Ware, Hertfordshire SG12 0DP (GB)

(74) Representative: **Teuten, Andrew John et al**
GlaxoSmithKline
Corporate Intellectual Property (CN9.25.1)
980 Great West Road
Brentford, Middlesex TW8 9GS (GB)

(56) References cited:
WO-A-91/11173 **WO-A-91/11496**
WO-A-92/00061 **WO-A-92/08447**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] This invention relates to aerosol formulations of use for the administration of medicaments by inhalation.

[0002] The use of aerosols to administer medicaments has been known for several decades. Such aerosols generally comprise the medicament, one or more chlorofluorocarbon propellants and either a surfactant or a solvent, such as ethanol. The most commonly used aerosol propellants for medicaments have been propellant 11 (CCl_3F) and/or propellant 114 ($\text{CF}_2\text{ClCF}_2\text{Cl}$) with propellant 12 (CCl_2F_2). However these propellants are now believed to provoke the degradation of stratospheric ozone and there is thus a need to provide aerosol formulations for medicaments which employ so called "ozone-friendly" propellants.

[0003] A class of propellants which are believed to have minimal ozone-depleting effects in comparison to conventional chlorofluorocarbons comprise fluorocarbons and hydrogen-containing chlorofluorocarbons, and a number of medicinal aerosol formulations using such propellant systems are disclosed in, for example, EP 0372777, WO91/04011, WO91/11173, WO91/11495 and WO91/14422. These applications are all concerned with the preparation of pressurised aerosols for the administration of medicaments and seek to overcome the problems associated with the use of the new class of propellants, in particular the problems of stability associated with the pharmaceutical formulations prepared. The applications all propose the addition of one or more of adjuvants such as alcohols, alkanes, dimethyl ether, surfactants (including fluorinated and non-fluorinated surfactants, carboxylic acids, polyethoxylates etc) and even conventional chlorofluorocarbon propellants in small amounts intended to minimise potential ozone damage.

[0004] Thus, for example EP 0372777 requires the use of 1,1,1,2-tetrafluoroethane in combination with both a cosolvent having greater polarity than 1,1,1,2-tetrafluoroethane (e.g. an alcohol or a lower alkane) and a surfactant in order to achieve a stable formulation of a medicament powder. In particular it is noted in the specification at page 3, line 7 that "it has been found that the use of propellant 134a (1,1,1,2-tetrafluoroethane) and drug as a binary mixture or in combination with a conventional surfactant such as sorbitan trioleate does not provide formulations having suitable properties for use with pressurised inhalers". Surfactants are generally recognised by those skilled in the art to be essential components of aerosol formulations, required not only to reduce aggregation of the medicament but also to lubricate the valve employed, thereby ensuring consistent reproducibility of valve actuation and accuracy of dose dispensed. Whilst WO91/11173, WO91/11495 and WO91/14422 are concerned with formulations comprising an admixture of drug and surfactant, WO91/04011 discloses medicinal aerosol formulations

in which the particulate medicaments are pre-coated with surfactant prior to dispersal in 1,1,1,2-tetrafluoroethane.

[0005] WO92/08447 discloses formulations comprising a drug which is surface coated with a surfactant and a hydrofluorocarbon propellant. WO91/11496 discloses use of HFA227 as an alternative propellant optionally in admixture with another propellant gas. US3,320,125 discloses CFC-containing pharmaceutical aerosol formulations in which use of sorbitan trioleate is avoided by use of propellant 11 at a concentration of 5-50%.

[0006] We have now surprisingly found that, in contradistinction to these teachings, it is in fact possible to obtain satisfactory dispersions of certain medicaments in fluorocarbon or hydrogen-containing chlorofluorocarbon propellants such as 1,1,1,2-tetrafluoroethane without recourse to the use of any surfactant or cosolvent in the composition, or the necessity to pre-treat the medicament prior to dispersal in the propellant.

[0007] There is thus provided in one aspect of the invention a pharmaceutical aerosol formulation for use in the administration of medicaments by inhalation characterised in that it consists essentially of particulate fluticasone propionate as medicament and 1,1,1,2-tetrafluoroethane as propellant and that the formulation is free of surfactant, wherein said medicament is present in an amount of 0.01% to 1% w/w relative to the total weight of the formulation. By "free of surfactant" is meant formulations which contain no significant amounts of surfactant, for example less than 0.0001% by weight of the medicament.

[0008] There is also provided a pharmaceutical aerosol formulation for use in the administration of medicaments by inhalation characterised in that it consists of particulate fluticasone propionate and 1,1,1,2-tetrafluoroethane wherein said fluticasone propionate is present in an amount of 0.01% to 1% w/w relative to the total weight of the formulation.

[0009] As another aspect the present invention provides use of a pharmaceutical aerosol formulation as hereinbefore defined for the manufacture of a medicament for the treatment of respiratory disorders.

[0010] The particle size of the particulate (e.g. micronised) medicament should be such as to permit inhalation of substantially all of the medicament into the lungs upon administration of the aerosol formulation and will thus be less than 100 microns, desirably less than 20 microns, and preferably in the range 1-10 microns, e.g. 1-5 microns.

[0011] The propellant for use in the invention is 1,1,1,2-tetrafluoroethane which has a sufficient vapour pressure to render it effective as a propellant.

[0012] It is desirable that the formulations of the invention contain no components which may provoke the degradation of stratospheric ozone. In particular it is desirable that the formulations are substantially free of chlorofluorocarbons such as CCl_3F , CCl_2F_2 and CF_3CCl_3 .

[0013] Further, formulations which are substantially free of volatile adjuvants such as a saturated hydrocarbon, for example propane, n-butane, isobutane, pentane and isopentane, or a dialkyl ether for example dimethyl ether, are preferred.

[0014] The formulations of the invention may be prepared by dispersal of the medicament in the propellant in an appropriate container, e.g. with the aid of sonication. The process is desirably carried out under anhydrous conditions to obviate any adverse effects of moisture on suspension stability.

[0015] The formulations according to the invention form weakly flocculated suspensions on standing but, surprisingly, these suspensions have been found to be easily redispersed by mild agitation to provide suspensions with excellent delivery characteristics suitable for use in pressurised inhalers, even after prolonged storage. Avoiding the use of formulation excipients e.g. surfactants, cosolvents etc in the aerosol formulations according to the invention is also advantageous since the formulations may be substantially taste and odour free, less irritant and less toxic than conventional formulations.

[0016] The chemical and physical stability and the pharmaceutical acceptability of the aerosol formulations according to the invention may be determined by techniques well known to those skilled in the art. Thus, for example, the chemical stability of the components may be determined by HPLC assay, for example, after prolonged storage of the product. Physical stability data may be gained from other conventional analytical techniques such as, for example, by leak testing, by valve delivery assay (average shot weights per actuation), by dose reproducibility assay (active ingredient per actuation) and spray distribution analysis.

[0017] The particle size distribution of the aerosol formulations according to the invention is particularly impressive and may be measured by conventional techniques, for example by cascade impaction or by the "Twin Impinger" analytical process. As used herein reference to the "Twin Impinger" assay means "Determination of the deposition of the emitted dose in pressurised inhalations using apparatus A" as defined in British Pharmacopoeia 1988, pages A204-207, Appendix XVII C. Such techniques enable the "respirable fraction" of the aerosol formulations to be calculated. As used herein reference to "respirable fraction" means the amount of active ingredient collected in the lower impingement chamber per actuation expressed as a percentage of the total amount of active ingredient delivered per actuation using the twin impinger method described above. The formulations according to the invention have been found to have a respirable fraction of 20% or more by weight of the medicament, preferably 25 to 70%, for example 30 to 60%.

[0018] The formulations for use in the invention may be filled into canisters suitable for delivering pharmaceutical aerosol formulations. Canisters generally com-

prise a container capable of withstanding the vapour pressure of the propellant such as a plastic or plastic-coated glass bottle or preferably a metal can, for example an aluminium can which may optionally be anodised, lacquer-coated and/or plastic-coated, which container is closed with a metering valve. The metering valves are designed to deliver a metered amount of the formulation per actuation and incorporate a gasket to prevent leakage of propellant through the valve. The gasket may comprise any suitable elastomeric material such as for example low density polyethylene, chlorobutyl, black and white butadiene-acrylonitrile rubbers, butyl rubber and neoprene. Suitable valves are commercially available from manufacturers well known in the aerosol industry, for example, from Valois, France (e.g. DF10, DF30, DF60), Bespak plc, UK (e.g. BK300, BK356) and 3M-Neotechnic Ltd, UK (e.g. Spraymiser™).

[0019] Conventional bulk manufacturing methods and machinery well known to those skilled in the art of pharmaceutical aerosol manufacture may be employed for the preparation of large scale batches for the commercial production of filled canisters. Thus, for example, in one bulk manufacturing method a metering valve is crimped onto an aluminium can to form an empty canister. The particulate medicament is added to a charge vessel and liquified propellant is pressure filled through the charge vessel into a manufacturing vessel. The drug suspension is mixed before recirculation to a filling machine and an aliquot of the drug suspension is then filled through the metering valve into the canister. Typically, in batches prepared for pharmaceutical use, each filled canister is check-weighed, coded with a batch number and packed into a tray for storage before release testing.

[0020] Each filled canister is conveniently fitted into a suitable channelling device prior to use to form a metered dose inhaler for administration of the medicament into the lungs or nasal cavity of a patient. Suitable channelling devices comprise for example a valve actuator and a cylindrical or cone-like passage through which medicament may be delivered from the filled canister via the metering valve to the nose or mouth of a patient e.g. a mouthpiece actuator. Metered dose inhalers are designed to deliver a fixed unit dosage of medicament per actuation or "puff", for example in the range of 10 to 5000 microgram medicament per puff.

[0021] Administration of medicament may be indicated for the treatment of mild, moderate or severe acute or chronic symptoms or for prophylactic treatment. It will be appreciated that the precise dose administered will depend on the age and condition of the patient and the frequency of administration and will ultimately be at the discretion of the attendant physician. Typically, administration may be one or more times, for example from 1 to 8 times per day, giving for example 1,2,3 or 4 puffs each time.

[0022] Suitable daily doses, may be, for example in the range 50 to 2000 microgram of fluticasone propionate depending on the severity of the disease.

[0023] Thus, for example, each valve actuation may deliver 25, 50, 125 or 250 microgram fluticasone propionate. Typically each filled canister for use in a metered dose inhaler contains 100, 160 or 240 metered doses or puffs of medicament.

[0024] Respiratory disorders such as, for example, asthma, can be treated by the administration by inhalation of an effective amount of a formulation as herein described.

[0025] The following non-limitative Examples serve to illustrate the invention.

Example 1

[0026] Micronised fluticasone propionate (24mg) was weighed into a clean, dry, plastic-coated glass bottle and 1,1,1,2-tetrafluoroethane (18.2g) was added from a vacuum flask. The bottle was quickly sealed with a blank aluminium ferrule. The resulting aerosol contained 0.132% w/w fluticasone propionate.

Examples 2 and 3

[0027] Micronised fluticasone propionate (66mg or 6.6mg) was weighed directly into each of 100 open aluminium cans and a metering valve was then crimped into place on each can. 1,1,1,2-Tetrafluoroethane (18.2g) was then added to each canister under pressure, through the valve, and each filled canister shaken to disperse the drug. The resulting inhalers contained 66 or 6.6mg fluticasone propionate and delivered 250 or 25 microgram fluticasone propionate per actuation (Examples 2 and 3 respectively).

Claims

1. A pharmaceutical aerosol formulation for use in the administration of medicaments by inhalation **characterised in that** it consists essentially of particulate fluticasone propionate as medicament, and 1,1,1,2-tetrafluoroethane as propellant, and that the formulation is free of surfactant, wherein said medicament is present in an amount of 0.01 to 1% w/w relative to the total weight of the formulation.
2. A pharmaceutical aerosol formulation for use in the administration of medicaments by inhalation **characterised in that** it consists of particulate fluticasone propionate and 1,1,1,2-tetrafluoroethane, wherein said fluticasone propionate is present in an amount of 0.01 to 1% w/w relative to the total weight of the formulation.
3. A pharmaceutical aerosol formulation according to claim 2 wherein fluticasone propionate is present in the formulation at a concentration of 0.132% w/w.

4. A pharmaceutical aerosol formulation according to claim 2 wherein fluticasone propionate is present in the formulation in an amount of 66mg per 18.2g of 1,1,1,2-tetrafluoroethane.
5. A pharmaceutical aerosol formulation according to claim 2 wherein fluticasone propionate is present in the formulation in an amount of 6.6mg per 18.2g of 1,1,1,2-tetrafluoroethane.
6. A pharmaceutical aerosol formulation according to any one of claims 1 to 5 wherein the use is use in the treatment of respiratory disorders.
7. A pharmaceutical aerosol formulation according to claim 6 wherein the respiratory disorder is asthma.
8. The use of a pharmaceutical aerosol formulation **characterised in that** it consists essentially of particulate fluticasone propionate as medicament, and 1,1,1,2-tetrafluoroethane as propellant, and that the formulation is free of surfactant, for the manufacture of a medicament for administration by inhalation, wherein said medicament is present in an amount of 0.01 to 1% w/w relative to the total weight of the formulation.
9. The use of a pharmaceutical aerosol formulation **characterised in that** it consists of particulate fluticasone propionate and 1,1,1,2-tetrafluoroethane, for the manufacture of a medicament for administration by inhalation, wherein said fluticasone propionate is present in an amount of 0.01 to 1% w/w relative to the total weight of the formulation.
10. Use according to claim 9 wherein fluticasone propionate is present in the formulation at a concentration of 0.132% w/w.
11. Use according to claim 9 wherein fluticasone propionate is present in the formulation in an amount of 66mg per 18.2g of 1,1,1,2-tetrafluoroethane.
12. Use according to claim 9 wherein fluticasone propionate is present in the formulation in an amount of 6.6mg per 18.2g of 1,1,1,2-tetrafluoroethane.
13. The use according to any one of claims 8 to 12 wherein the use is use in the treatment of respiratory disorders.
14. The use according to claim 13 wherein the respiratory disorder is asthma.
15. A canister comprising a container closed with a metering valve which contains a pharmaceutical aerosol formulation according to any one of claims 1 to 7.

16. A canister according to claim 15 wherein the container is an aluminium can.
17. A canister according to claim 16 wherein the aluminium can is anodised, lacquer coated and/or plastic coated.
18. A process for preparing a pharmaceutical aerosol formulation for use in the administration of medicaments by inhalation **characterised in that** it consists essentially of particulate fluticasone propionate as medicament, and 1,1,1,2-tetrafluoroethane as propellant, and that the formulation is free of surfactant, wherein said medicament is present in an amount of 0.01-1% w/w relative to the total weight of the formulation, which process comprises dispersing the medicament in the propellant.
19. A process for preparing a pharmaceutical aerosol formulation for use in the administration of medicaments by inhalation **characterised in that** it consists of particulate fluticasone propionate and 1,1,1,2-tetrafluoroethane, wherein said fluticasone propionate is present in an amount of 0.01-1% w/w relative to the total weight of the formulation, which process comprises dispersing the fluticasone propionate in the 1,1,1,2-tetrafluoroethane.
20. A process according to claim 19 wherein fluticasone propionate is present in the formulation at a concentration of 0.132% w/w.
21. A process according to claim 19 wherein fluticasone propionate is present in the formulation in an amount of 66mg per 18.2g of 1,1,1,2-tetrafluoroethane.
22. A process according to claim 19 wherein fluticasone propionate is present in the formulation in an amount of 6.6mg per 18.2g of 1,1,1,2-tetrafluoroethane.

Patentansprüche

1. Pharmazeutische Aerosolformulierung zur Verwendung in der Verabreichung von Medikamenten durch Inhalation, **dadurch gekennzeichnet, daß** sie im wesentlichen aus teilchenförmigem Fluticasonpropionat als Medikament und 1,1,1,2-Tetrafluorethan als Treibmittel besteht, und daß die Formulierung frei von Tensid ist, worin das Medikament in einer Menge von 0,01 bis 1 % G/G relativ zum Gesamtgewicht der Formulierung vorhanden ist.
2. Pharmazeutische Aerosolformulierung zur Verwendung in der Verabreichung von Medikamenten durch Inhalation, **dadurch gekennzeichnet, daß**

sie aus teilchenförmigem Fluticasonpropionat und 1,1,1,2-Tetrafluorethan besteht, worin das Fluticasonpropionat in einer Menge von 0,01 bis 1 % G/G relativ zum Gesamtgewicht der Formulierung

3. Pharmazeutische Aerosolformulierung gemäß Anspruch 2, worin Fluticasonpropionat in der Formulierung in einer Konzentration von 0,132 % G/G vorhanden ist.
4. Pharmazeutische Aerosolformulierung gemäß Anspruch 2, worin Fluticasonpropionat in der Formulierung in einer Menge von 66 mg auf 18,2 g 1,1,1,2-Tetrafluorethan vorhanden ist.
5. Pharmazeutische Aerosolformulierung gemäß Anspruch 2, worin Fluticasonpropionat in der Formulierung in einer Menge von 6,6 mg auf 18,2 g 1,1,1,2-Tetrafluorethan vorhanden ist.
6. Pharmazeutische Aerosolformulierung gemäß einem der Ansprüche 1 bis 5, worin die Verwendung die Verwendung in der Behandlung von Atemwegsstörungen ist.
7. Pharmazeutische Aerosolformulierung gemäß Anspruch 6, worin die Atemwegsstörung Asthma ist.
8. Verwendung einer pharmazeutischen Aerosolformulierung, **dadurch gekennzeichnet, daß** sie im wesentlichen aus teilchenförmigem Fluticasonpropionat als Medikament und 1,1,1,2-Tetrafluorethan als Treibmittel besteht, und daß die Formulierung frei von Tensid ist, zur Herstellung eines Medikaments zur Verabreichung durch Inhalation, worin das Medikament in einer Menge von 0,01 bis 1 % G/G relativ zum Gesamtgewicht der Formulierung vorhanden ist.
9. Verwendung einer pharmazeutischen Aerosolformulierung, **dadurch gekennzeichnet, daß** sie aus teilchenförmigem Fluticasonpropionat und 1,1,1,2-Tetrafluorethan besteht, zur Herstellung eines Medikaments zur Verabreichung durch Inhalation, worin das Fluticasonpropionat in einer Menge von 0,01 bis 1 % G/G relativ zum Gesamtgewicht der Formulierung vorhanden ist.
10. Verwendung gemäß Anspruch 9, worin Fluticasonpropionat in der Formulierung in einer Konzentration von 0,132 % G/G vorhanden ist.
11. Verwendung gemäß Anspruch 9, worin Fluticasonpropionat in der Formulierung in einer Menge von 66 mg auf 18,2 g 1,1,1,2-Tetrafluorethan vorhanden ist.
12. Verwendung gemäß Anspruch 9, worin Fluticason-

propionat in der Formulierung in einer Menge von 6,6 mg auf 18,2 g 1,1,1,2-Tetrafluorethan vorhanden ist.

13. Verwendung gemäß einem der Ansprüche 8 bis 12, worin die Verwendung die Verwendung in der Behandlung von Atemwegsstörungen ist.
14. Verwendung gemäß Anspruch 13, worin die Atemwegsstörung Asthma ist.
15. Kanister, umfassend einen mit einem Dosierventil verschlossenen Behälter, der eine pharmazeutische Aerosolformulierung gemäß einem der Ansprüche 1 bis 7 enthält.
16. Kanister gemäß Anspruch 15, worin der Behälter eine Aluminiumdose ist.
17. Kanister gemäß Anspruch 16, worin das Aluminium eloxiert, lackiert und/oder kunststoffbeschichtet ist.
18. Verfahren zur Herstellung einer pharmazeutischen Aerosolformulierung zur Verwendung in der Verabreichung von Medikamenten durch Inhalation, **dadurch gekennzeichnet, daß** sie im wesentlichen aus teilchenförmigem Fluticasonpropionat als Medikament und 1,1,1,2-Tetrafluorethan als Treibmittel besteht, und daß die Formulierung frei von Tensid ist, worin das Medikament in einer Menge von 0,01 bis 1 % G/G relativ zum Gesamtgewicht der Formulierung vorhanden ist, wobei das Verfahren das Dispergieren des Medikaments im Treibmittel umfaßt.
19. Verfahren zur Herstellung einer pharmazeutischen Aerosolformulierung zur Verwendung in der Verabreichung von Medikamenten durch Inhalation, **dadurch gekennzeichnet, daß** sie aus teilchenförmigem Fluticasonpropionat und 1,1,1,2-Tetrafluorethan besteht, worin das Fluticasonpropionat in einer Menge von 0,01 bis 1 % G/G relativ zum Gesamtgewicht der Formulierung vorhanden ist, wobei das Verfahren das Dispergieren des Fluticasonpropionats im 1,1,1,2-Tetrafluorethan umfaßt.
20. Verfahren gemäß Anspruch 19, worin Fluticasonpropionat in der Formulierung in einer Konzentration von 0,132 % G/G vorhanden ist.
21. Verfahren gemäß Anspruch 19, worin Fluticasonpropionat in der Formulierung in einer Menge von 66 mg auf 18,2 g 1,1,1,2-Tetrafluorethan vorhanden ist.
22. Verfahren gemäß Anspruch 19, worin Fluticasonpropionat in der Formulierung in einer Menge von 6,6 mg auf 18,2 g 1,1,1,2-Tetrafluorethan vorhan-

den ist.

Revendications

1. Formulation pharmaceutique sous forme d'aérosol utile dans l'administration de médicaments par inhalation, **caractérisée en ce qu'elle** consiste essentiellement en propionate de fluticasone particulaire en tant que médicament, et en 1,1,1,2-tétrafluoroéthane en tant qu'agent propulseur, et que la formulation est dépourvue de tensioactif, dans laquelle ledit médicament est présent en une quantité de 0,01 à 1% en poids/poids par rapport au poids total de la formulation.
2. Formulation pharmaceutique sous forme d'aérosol utile dans l'administration de médicaments par inhalation, **caractérisée en ce qu'elle** consiste en propionate de fluticasone particulaire et en 1,1,1,2-tétrafluoroéthane, dans laquelle ledit propionate de fluticasone est présent en une quantité de 0,01 à 1% en poids/poids par rapport au poids total de la formulation.
3. Formulation pharmaceutique sous forme d'aérosol suivant la revendication 2, dans laquelle le propionate de fluticasone est présent dans la formulation à une concentration de 0,132% en poids/poids.
4. Formulation pharmaceutique sous forme d'aérosol suivant la revendication 2, dans laquelle le propionate de fluticasone est présent dans la formulation en une quantité de 66 mg pour 18,2 g de 1,1,1,2-tétrafluoroéthane.
5. Formulation pharmaceutique sous forme d'aérosol suivant la revendication 2, dans laquelle le propionate de fluticasone est présent dans la formulation en une quantité de 6,6 mg pour 18,2 g de 1,1,1,2-tétrafluoroéthane.
6. Formulation pharmaceutique sous forme d'aérosol suivant l'une quelconque des revendications 1 à 5, dans laquelle l'utilisation est une utilisation dans le traitement de troubles respiratoires.
7. Formulation pharmaceutique sous forme d'aérosol suivant la revendication 6, dans laquelle le trouble respiratoire est l'asthme.
8. Utilisation d'une formulation pharmaceutique sous forme d'aérosol, **caractérisée en ce qu'elle** consiste essentiellement en propionate de fluticasone particulaire en tant que médicament, et en 1,1,1,2-tétrafluoroéthane en tant qu'agent propulseur, et que la formulation est dépourvue de tensioactif, pour la fabrication d'un médicament pour

l'administration par inhalation, dans laquelle ledit médicament est présent en une quantité de 0,01 à 1% en poids/poids par rapport au poids total de la formulation.

9. Utilisation d'une formulation pharmaceutique sous forme d'aérosol, **caractérisée en ce qu'elle** consiste en propionate de fluticasone particulaire et en 1,1,1,2-tétrafluoroéthane pour la fabrication d'un médicament pour l'administration par inhalation, dans laquelle ledit propionate de fluticasone est présent en une quantité de 0,01 à 1% en poids/poids par rapport au poids total de la formulation.

10. Utilisation suivant la revendication 9, dans laquelle le propionate de fluticasone est présent dans la formulation à une concentration de 0,132% en poids/poids.

11. Utilisation suivant la revendication 9, dans laquelle le propionate de fluticasone est présent dans la formulation en une quantité de 66 mg pour 18,2 g de 1,1,1,2-tétrafluoroéthane

12. Utilisation suivant la revendication 9, dans laquelle le propionate de fluticasone est présent dans la formulation en une quantité de 6,6 mg pour 18,2 g de 1,1,1,2-tétrafluoroéthane.

13. Utilisation suivant l'une quelconque des revendications 8 à 12, dans laquelle l'utilisation est une utilisation dans le traitement de troubles respiratoires.

14. Utilisation suivant la revendication 13, dans laquelle le trouble respiratoire est l'asthme.

15. Récipient comprenant un réservoir fermé par une vanne doseuse qui contient une formulation pharmaceutique sous forme d'aérosol suivant l'une quelconque des revendications 1 à 7.

16. Récipient suivant la revendication 15, dans laquelle le réservoir est une boîte en aluminium.

17. Récipient suivant la revendication 16, dans laquelle la boîte d'aluminium est anodisée, revêtue de laque et/ou revêtue de matière plastique.

18. Procédé pour la préparation d'une formulation pharmaceutique sous forme d'aérosol utile dans l'administration de médicaments par inhalation, **caractérisé en ce qu'elle** consiste essentiellement en propionate de fluticasone en tant que médicament, et en 1,1,1,2-tétrafluoroéthane en tant qu'agent propulseur, et que la formulation est dépourvue de tensioactif, dans laquelle ledit médicament est présent en une quantité de 0,01 à 1% -en poids/poids par rapport au poids total de la formulation, lequel pro-

cédé comprend la dispersion du médicament dans l'agent propulseur.

19. Procédé pour la préparation d'une formulation pharmaceutique sous forme d'aérosol utile dans l'administration de médicaments par inhalation, **caractérisé en ce qu'elle** consiste en propionate de fluticasone particulaire et en 1,1,1,2-tétrafluoroéthane, dans laquelle ledit propionate de fluticasone est présent en une quantité de 0,01 à 1% en poids/poids par rapport au poids total de la formulation, lequel procédé comprend la dispersion du propionate de fluticasone dans le 1,1,1,2-tétrafluoroéthane.

20. Procédé suivant la revendication 19, dans lequel le propionate de fluticasone est présent dans la formulation à une concentration de 0,132% en poids/poids.

21. Procédé suivant la revendication 19, dans lequel le propionate de fluticasone est présent dans la formulation en une quantité de 66 mg pour 18,2 g de 1,1,1,2-tétrafluoroéthane.

22. Procédé suivant la revendication 19, dans lequel le propionate de fluticasone est présent dans la formulation en une quantité de 6,6 mg pour 18,2 g de 1,1,1,2-tétrafluoroéthane.